

# Journal of Fish Biology

## Understanding and managing fish populations: keeping the toolbox fit for purpose --Manuscript Draft--

<b>Manuscript Number:</b>	SP 17-012R2
<b>Full Title:</b>	Understanding and managing fish populations: keeping the toolbox fit for purpose
<b>Short Title:</b>	Tools for understanding fish populations
<b>Article Type:</b>	FSBI Symposium SI Review Paper
<b>Keywords:</b>	archaeology, genetics, modelling, surveys, stable isotopes, telemetry
<b>Corresponding Author:</b>	Josephine Paris University of Exeter Exeter, UNITED KINGDOM
<b>Corresponding Author Secondary Information:</b>	
<b>Corresponding Author's Institution:</b>	University of Exeter
<b>Corresponding Author's Secondary Institution:</b>	
<b>First Author:</b>	Josephine Paris
<b>First Author Secondary Information:</b>	
<b>Order of Authors:</b>	Josephine Paris Krista Sherman Brian Hayden, PhD Jamie Stevens, PhD
<b>Order of Authors Secondary Information:</b>	
<b>Manuscript Region of Origin:</b>	UNITED KINGDOM
<b>Abstract:</b>	Wild fish populations are currently experiencing unprecedented pressures, which are projected to intensify in the coming decades. Developing a thorough understanding of the influences of both biotic and abiotic factors on fish populations is a salient issue in contemporary fish conservation and management. During the 50th Anniversary of the Fisheries Society of the British Isles, University of Exeter, 2017, scientists from diverse research backgrounds gathered to discuss key topics under the broad umbrella of 'Understanding Fish Populations'. Below, the output of one such discussion group is detailed, focusing on tools used to investigate natural fish populations. Five main groups of approaches were identified: (i) Tagging and telemetry; (ii) Molecular tools; (iii) Survey tools; (iv) Statistical and modelling tools; and (v) Tissue analyses. The appraisal covered current challenges and potential solutions for each of these topics. In addition, three key themes were identified as applicable across all tool-based applications. These included data management, public engagement, and fisheries policy and governance. The continued innovation of tools and capacity to integrate interdisciplinary approaches into the future assessment and management of fish populations is highlighted as an important focus for the next 50 years of fisheries research.

**[Type text] [Type text] 14/07/15 Ethics Questionnaire for JFB**

***Submitted manuscripts will only be considered if the experimental methods employed are ethically justified. Please answer all questions. If you have answered 'yes' to questions 4 to 7, you should include an Ethics paragraph in the Methods section of your manuscript which justifies your methods used. You should complete this questionnaire based on all fishes used in your experiment. For example, if you used live fishes as prey in predation experiments, this is a lethal endpoint for the prey fish (see Questions 5 & 6). Please read the Editorial published in JFB 68, 1-2, for full information on JFB ethics. PLEASE SUBMIT THE COMPLETED QUESTIONNAIRE WITH YOUR MANUSCRIPT ONLINE THROUGH EDITORIAL MANAGER.***

***Corresponding author's name: Dr Jamie Stevens***

***Question 1: Were fishes collected as part of faunal surveys? NO***

***Question 2: If you have undertaken experimental work, has the care and use of experimental animals complied with local and or national animal welfare laws, guidelines and policies? NO***

***If 'Yes', state these and provide suitable evidence (e.g. for the U.K. a Home Office PPL number is sufficient), both here and in the manuscript, that protocols have undergone an ethical review process by an institutional animal care and use (or similar) committee, a local ethics committee, or by appropriately qualified scientific and lay colleagues.***

***[Type text] [Type text] 14/07/15***

***If 'No', because these laws do not exist in your country, please state this. Alternatively, if you carried out purely observational work so ethical permission was not considered necessary please state this both here and in the manuscript.***

***Purely observational work***

***Question 3: Were fishes killed during or at the end of your experiment (e.g. for tissue sampling)? NO***

***If 'Yes', what method was used? Please provide details both here and in the manuscript.***

***Question 4: Have you performed surgical procedures? NO If 'Yes', please give brief details of the surgery here. Full details should be given in the manuscript. If the procedures caused more than slight pain or distress, did you use appropriate sedation, analgesia and anaesthesia, with post-operative care? Please provide full details and justification both here and within the manuscript including type and concentration of anaesthetic.***

**[Type text] [Type text] 14/07/15**

**Question 5: *Did you use experimental conditions that severely distressed any fishes involved in your experiments? NO***

***If 'Yes', state the conditions and how they can be justified. What humane endpoints were used to minimise the effects? Please provide full justification within the methods section of your manuscript.***

**Question 6: *Did any of the experimental procedures, particularly those that involve lethal endpoints (e.g. predation studies, toxicity testing), cause lasting harm to sentient fishes? NO***

***If 'Yes', provide details both here and in the methods section of your manuscript. Normally these procedures will be considered unacceptable by JFB unless any harm caused can be justified against the benefits gained.***

**Question 7: *Did any of your procedures involve sentient, un-anaesthetised animals paralysed by chemical agents such as muscle relaxants? NO***

***If 'Yes', provide details both here and in the methods section of your manuscript. Normally these procedures will be considered unacceptable by JFB.***



Received 31 July 2017

Accepted 20 Decemebr 2017

# Understanding and managing fish populations: keeping the toolbox fit for purpose

J. R. PARIS<sup>1\*</sup>, K. D. SHERMAN<sup>1\*</sup>, E. BELL<sup>2</sup>, C. BOULENGER<sup>3</sup>, C. DELORD<sup>3</sup>, M. B. M. EL-MAHDI<sup>4</sup>, E. A. FAIRFIELD<sup>2</sup>, A. M. GRIFFITHS<sup>1</sup>, C. GUTMANN ROBERTS<sup>5</sup>, R. D. HEDGER<sup>6</sup>, L. E. HOLMAN<sup>7</sup>, L. H. HOOPER<sup>1</sup>, N. E. HUMPHRIES<sup>8</sup>, I. KATSIADAKI<sup>9</sup>, R. A. KING<sup>1</sup>, A. LEMOPOULOS<sup>10</sup>, C. J. PAYNE<sup>11</sup>, G. PEIRSON<sup>12</sup>, K. K. RICHTER<sup>13</sup>, M. I. TAYLOR<sup>2</sup>, C. N. TRUEMAN<sup>7</sup>, B. HAYDEN<sup>14†</sup> & J. R. STEVENS<sup>1†#</sup>

<sup>1</sup>Biosciences, University of Exeter, Stocker Road, Exeter EX4 4QD, U.K., <sup>2</sup>School of Biological Science, University of East Anglia, Norwich Research Park, Norwich NR4 7UG, U.K., <sup>3</sup>ESE, Ecology and Ecosystem Health, Agrocampus Ouest, INRA, 35042 Rennes, France, <sup>4</sup>Faculty of Science, Laboratory of Molecular Genetics and Molecular Biology, Zoology Department, South Valley University, Qena 83523, Egypt, <sup>5</sup>Department of Life and Environmental Sciences, Faculty of Science and Technology, Bournemouth University, Bournemouth BH12 5BB, U.K., <sup>6</sup>Norwegian Institute for Nature Research, Trondheim, NO-7485, Norway, <sup>7</sup>Ocean and Earth Science, National Oceanography Centre, University of Southampton, European Way, Southampton SO14 3ZH, U.K., <sup>8</sup>Marine Biological Association of the United Kingdom, The Laboratory, Citadel Hill, Plymouth PL1 2PB, U.K., <sup>9</sup>Cefas, Weymouth Laboratory, Weymouth, DT4 8UB, U.K., <sup>10a</sup>Department of Environmental and Biological Sciences, University of Eastern Finland, P.O. Box 111, FI-80101 Joensuu, Finland, <sup>10b</sup>Department of Biology, University of Turku, FI- 20014, Turku, Finland, <sup>11</sup>Institute

of Aquaculture, Faculty of Natural Sciences, University of Stirling, Stirling FK9 4LA, U.K.,

<sup>12</sup>Environment Agency, Research Analysis and Evaluation, Mance House, Kidderminster

DY11 7RA, U.K., <sup>13</sup>BioArCh, Department of Archaeology, University of York, York YO10

5DD, U.K. and <sup>14</sup>Biology Department, Canadian Rivers Institute, University of New

Brunswick, Fredericton, NB, E3B 5A3, Canada

**Running title:** TOOLS FOR UNDERSTANDING FISH POPULATIONS

#Author to whom correspondence should be addressed. Tel.: +44 1392 723 775; email:

j.r.stevens@exeter.ac.uk

\* Joint first authors; † Joint senior authors

Wild fish populations are currently experiencing unprecedented pressures, which are projected to intensify in the coming decades. Developing a thorough understanding of the influences of both biotic and abiotic factors on fish populations is a salient issue in contemporary fish conservation and management. During the 50th Anniversary Symposium of The Fisheries Society of the British Isles at the University of Exeter, UK, in July 2017, scientists from diverse research backgrounds gathered to discuss key topics under the broad umbrella of ‘Understanding Fish Populations’. Below, the output of one such discussion group is detailed, focusing on tools used to investigate natural fish populations. Five main groups of approaches were identified: tagging and telemetry; molecular tools; survey tools; statistical and modelling tools; tissue analyses. The appraisal covered current challenges and potential solutions for each of these topics. In addition, three key themes were identified as applicable across all tool-based applications. These included data management, public engagement and fisheries policy and governance. The continued innovation of tools and capacity to integrate interdisciplinary approaches into the future assessment and management of fish populations is highlighted as an important focus for the next 50 years of fisheries research.

Key words: archaeology, genetics, modelling, surveys, stable isotopes, telemetry



## INTRODUCTION

Approximately 30% of fish species have been overexploited (FAO, 2014), representing significant losses to biodiversity, ecosystem services and socioeconomic contributions (Worm *et al.*, 2009). In light of the increasing challenges presented by climate change and other natural and anthropogenic stressors (Gordon *et al.*, 2018), an improved understanding of fish populations is critical to facilitate effective management and conservation initiatives. During July 2017, The Fisheries Society of the British Isles held its 50th Anniversary Symposium at the University of Exeter, UK, under the broad umbrella of ‘Understanding Fish Populations’. To highlight key knowledge gaps and opportunities, we report the outcome of a working group convened at the symposium, which was tasked with considering the theme of tools for understanding fish populations. The scope of the discussion spanned diverse areas including spatial ecology and migration patterns, genetics and evolutionary biology, physiology, trophic ecology and developmental and population biology. In this article, we consider major advances in the use of tools across broad areas of fish biology and identify knowledge gaps and potential solutions in each area in order to guide and inform future research and to better understand and protect wild fish populations.

## TAGGING AND TELEMETRY

A significant problem hampering the study of fish, marine benthic species in particular, is that of determining their geographical locations at fine scales, over long durations. Tagging and telemetry involves the application of external and or internal tags or devices to manually or passively track fish movement (Cooke *et al.*, 2013). Both forms can be particularly challenging in the marine environment, though manual tracking can work well at feeding grounds and at spawning aggregations (Murchie *et al.*, 2015), while passive

tracking has valuable applications along known migration routes (Dahlgren *et al.*, 2016), for example, as anadromous and catadromous species migrate in and out of river estuaries (Lauridsen *et al.*, 2017). Suites of tools exist for such tasks [*e.g.* acoustic transmitters, passive information transponder (PIT) and Floy tags, radio, archival, *etc.*] and have been routinely used to understand the spatial ecology of a range of fish taxa (Bograd *et al.*, 2010). With technological improvements in tags and tracking equipment, the field has grown vastly in recent decades (Pine *et al.*, 2003; Jepsen *et al.*, 2015). We briefly highlight some of the tags and telemetry options commonly used by researchers along with a discussion of some of the limitations and challenges associated with these tools.

Archival data storage tags (DST), which can collect data on both the internal and external environments of fish are the only method available to assess internal states (*e.g.* bioenergetics; Cooke *et al.*, 2016). DSTs, however, currently only provide information on the environment experienced by the tagged fish if the tag is recovered, meaning these data are lost if recapture rates are low as is often the case in fish tagging surveys. Communication history acoustic tags (CHAT), which transmit data to nearby transponder receivers are a promising alternative. Since there have been relatively few uses of this tag type (Voegeli *et al.*, 2001; Hight & Lowe, 2007), there is potential for development in this area. Pop-off DSTs are also becoming available and will no doubt prove very useful once problems associated with size and recoverability are resolved.

Pop-up Satellite Archival Tags (PSAT), which detach from the tagged fish after some time at sea and transmit telemetry data to overpassing satellites, are currently limited in terms of hardware, software and satellite reception. PSATs are large, so are limited in use for larger, often highly migratory individuals and may also affect fish behaviour (Methling *et al.*, 2011). Additionally, battery failure, antenna damage, or mechanical failure may limit registration or transmission of data (Hays *et al.*, 2007; Musyl *et al.*, 2011). PSAT technology

is relatively new, so future reductions in size and weight and also improvement in reliability can be expected. In terms of software, PSATs currently only transmit limited amounts of data due to transmission costs and the short time that the receiving satellite is above the horizon. Future software development is required to reduce transmission costs, optimize data transmission and provide more flexibility for users to tailor controls, in order to provide higher resolution data at the desired temporal scale. An increase in the number of satellite platforms that can receive PSAT data would help to improve reception issues. Interference on frequencies selected for tags at certain geographical locations (see Musyl *et al.*, 2011) also requires consideration.

Acoustic telemetry offers autonomous, continuous monitoring (Heupel *et al.*, 2006) and has the potential to significantly enhance our understanding of marine predator habitat use, activity patterns and resource partitioning (Hussey *et al.*, 2015). Acoustic arrays have been used in many studies elucidating fish movements ( Papastamatiou *et al.*, 2013; Lea *et al.*, 2016) and transmitters have been used more innovatively to measure trophic interactions (Halfyard *et al.*, 2017). Issues remain however, in the significant cost and effort involved in deploying and maintaining acoustic arrays.

Organisations such as the Ocean Tracking Network (OTN; [oceantrackingnetwork.org](http://oceantrackingnetwork.org); Whoriskey *et al.*, 2015) and the Integrated Marine Observing System (IMOS) animal tracking database Australian Animal Tracking Network ([www.imos.org.au/facilities/animaltracking](http://www.imos.org.au/facilities/animaltracking)) both maintain acoustic infra-structure in the form of deployed receivers (arrays or curtains) in key ecological areas into which researchers are free to release tagged animals. These initiatives substantially reduce the cost and risk associated with acoustic tracking projects and similar approaches can be applied globally (*e.g.* a European tracking network is currently being developed). Furthermore, integration of standardized data repositories along with a comprehensive set of analytical tools to ensure rapid and sophisticated analysis of acoustic

array data (Lea *et al.*, 2016) would lead to new insights into the spatial ecology of fish. Further technological developments such as the use of autonomous underwater vehicles (AUV) to perform routine data download operations, or even complement fixed acoustic receivers (Davis *et al.*, 2016), will make acoustic telemetry increasingly affordable and accessible to more researchers. Continued collaborations with established regional and international tracking networks, together with the ever-increasing sophistication, miniaturisation, durability and cost reduction of tags promises an increasingly important role for acoustic telemetry in our understanding of fish ecology.

## MOLECULAR TOOLS

### POPULATION GENETICS AND GENOMICS

Using genetic tools to understand fish genetic diversity and population structure has wide-ranging applications for evolutionary biology and the conservation and management of fish stocks. Until recently, molecular techniques such as mitochondrial sequencing and the analysis of microsatellite loci have been used most commonly to explore intra-specific variation in fish and many other organisms (Ferguson & Danzmann, 1998; Chistiakov *et al.*, 2006). More recently, however, the increased availability and cost efficiency of high-throughput sequencing, which is capable of producing millions of sequencing reads [*e.g.* RADseq (Davey & Blaxter, 2011) and RNAseq (Illumina; [www.illumina.com](http://www.illumina.com))], has revolutionized the fields of population and conservation genetics (Allendorf *et al.* 2010). It is, however, important to understand what extra information high-throughput sequencing data can provide, the biases involved in study design and data generation and also how its usage

might be optimized. Here, we seek to identify knowledge gaps in the field of fish population genetics and contemplate how this area of research may evolve in the future.

Attaining high quality, clean DNA for large numbers of individuals is paramount for downstream sequencing processes, but in some cases can be challenging. Biological samples can often be compromised during sampling or transport, potentially rendering field efforts futile. Population genetic studies on fish frequently require sampling from river transects or remote locations at sea and so portable laboratories for sampling, storing and extracting DNA would be welcomed. At the same time, emerging technologies, *e.g.* the MinION USB sequencer ([www.nanoporetech.com/products/minion](http://www.nanoporetech.com/products/minion)), have the potential to revolutionize when and where genetic data can be generated. Most new technologies are currently restricted to sequencing small genomes, such as those of bacteria, but with on-going improvements, these technologies open up the possibility of being able to sequence DNA in real-time in the field (Hayden, 2015). Recently, the MinION technology has started to be used in hybrid assemblies with Illumina short reads (Austin *et al.*, 2017) and *de novo* eukaryotic genomes (including fish) are in progress (Jansen *et al.*, 2017).

Alongside population genetic studies, research based on whole genome data is emerging and the genomes of several commercially important species have now been published (*e.g.* Atlantic cod *Gadus morhua* L. 1758; Star *et al.*, 2011; Atlantic salmon *Salmo salar* L. 1758; Lien *et al.*, 2016). While the ever-reducing cost of whole genome sequencing provides opportunities to sequence and publish more fish genomes, but, in our view, the key priority is not simply publishing genomes, it is also high-quality genome annotation. Gene annotation and accurate knowledge of the function of different identified regions is of extreme importance if genomic tools are to be used reliably in conservation and management (Ekblom & Wolf, 2014). Therefore, projects such as the ‘Functional Annotation of All Salmonid Genomes’ (Macqueen *et al.*, 2017) should be encouraged and developed. It is also

important not to underestimate or neglect the computing power and bioinformatics expertise required to produce high quality genome scaffolds and annotations and also to recognize and account for biases in next generation sequencing data (Benestan *et al.*, 2017).

Furthermore, population genetic approaches are usually focused on a single species. Consequently, there is a mismatch between studies of a single species genotyped at high resolution, but generally at small spatial scales (*e.g.* population genetics, often using hundreds to thousands of markers through genotyping by sequencing (GBS) or genome-wide association studies (GWAS)) and studies of multiple species at larger spatial scales but using lower resolution markers [*e.g.* phylogeography or biodiversity assessments using metabarcoding or mitochondrial (mt)DNA sequencing]. Nonetheless, the widespread application of molecular resources has led to the accumulation of rich datasets across a broad range of species, geographical regions and time periods (Blanchet *et al.*, 2017). Accordingly, we anticipate that this aggregation of data may allow the underlying processes that drive genetic variability across these regions and times to be revealed, enabling a broader testing of theories in population genetics and evolution (Ellegren & Galtier, 2016; Pauls *et al.*, 2014).

Such studies will require the combination of high genetic resolution markers across large spatial scales, which is a non-trivial task, especially when dealing with non-model species. Three challenges arise in such cases: firstly, the financial investment required to obtain reliable datasets for several species remains significant. Despite reductions in sequencing costs, it may be financially sensible to rely on more classical markers such as microsatellites or small subsets of single nucleotide polymorphisms (SNP). Secondly, there is a need for a standardized framework in order to make datasets comparable across different species and regions. This standardization must occur when collecting samples, characterizing markers ( Ellis *et al.*, 2011; Helyar *et al.* 2011) and during the subsequent data analysis to streamline user choices (Paris *et al.* 2017), which may bias the biological interpretation of

data (Rodríguez-Ezpeleta *et al.*, 2016). It is therefore important that researchers use common methods to isolate and characterize markers for entire sets of focal species and provide full access to detailed analyses when datasets are generated.

Finally, as multi-species approaches remain scarce, there is a need to define hypotheses at the beginning of such investigations. In this respect, simulation tools (Laval & Excoffier, 2004; Peng & Kimmal, 2005; Neuenschwander, 2006) are particularly useful for testing complex hypotheses and also for predictive purposes. Moreover, the integration of mathematical and statistical models with fish population genetics would be useful for revealing genotype-phenotype interactions (Ritchie *et al.*, 2015), evolutionary signatures (Stark *et al.*, 2007), functional DNA elements (Schridder & Kern, 2014), spatial dynamics (Guillot *et al.* 2009) and species-genetic diversity correlations (SGDC; Vellend 2003; Vellend *et al.*, 2014).

## ENVIRONMENTAL DNA

The use of environmental (e)DNA to identify the presence and understand the distribution of fish has expanded rapidly in the past decade. eDNA is a polydisperse mixture (Turner *et al.*, 2014; Wilcox *et al.*, 2015) of various biological material ranging from entire cellular fragments to extracellular DNA, which is isolated from environmental samples such as water or sediment. Such techniques are used for species identification and food security purposes. Universal primers that target mtDNA can be applied for identifying species presence (Yamamoto *et al.*, 2016) or to gain information about species interactions (*e.g.* food-web construction; Sousa *et al.* 2016).

An important component of this work is validating the results from eDNA surveys with traditional fish-survey methods. In both freshwater and marine environments, eDNA has

compared favourably with traditional fish survey methods (Thomsen *et al.*, 2012; Hänfling *et al.*, 2016). eDNA, however, was found to be less effective compared with experienced snorkel surveys (Ulibarri *et al.*, 2017). This underpins the importance of validation with traditional techniques, especially in spatially heterogeneous and complex aquatic environments (Shogren *et al.*, 2017).

The development of effective PCR primers is central to the successful application of eDNA (Freeland, 2016; MacDonald & Sarre, 2017). As a result, a vast range of primer sets are available for fishes (Doi *et al.*, 2015; Clusa *et al.*, 2017). Metabarcoding primers, that simultaneously amplify eDNA from many fish species, have also been developed for monitoring entire fish communities (Miya *et al.*, 2015; Valentini *et al.*, 2016).

Beyond inferring if a fish species is present in the sampled location, researchers have begun to investigate if eDNA can provide further information regarding fish populations. The use of eDNA to infer population-level variation has been demonstrated (Uchii *et al.*, 2015; Sigsgaard *et al.*, 2016), but is still in its infancy. Similarly, although attempts to link eDNA concentration and fish biomass have shown promising results (Lacoursière-Rousse *et al.*, 2016; Yamamoto *et al.*, 2016), but further development is required to improve the accuracy of these measurements. For techniques utilising eDNA to be optimized, however, preexisting molecular information needs to be accessible. A number of publicly available databases [*e.g.* NCBI Genbank ([www.ncbi.nlm.nih.gov/genbank/](http://www.ncbi.nlm.nih.gov/genbank/)) and BOLD ([www.boldsystems.org](http://www.boldsystems.org))] hold a vast array of molecular data, but there is still a need for further mitochondrial genome sequencing to allow optimal usage of molecular identification techniques.

## MICROBIOMES



Analysis of a microbiome can provide novel insights into the health and biology of fish populations. Traditional culture-dependent tools used to map the commensal microbiota community in teleost fish are often time-consuming, expensive and subjected to bias as only 0.1–10% of bacteria can be cultured *in vitro* (Amann *et al.* 1995; Austin, 2006). More recently, rapid culture-independent tools such as 16S ribosomal (r)RNA targeted sequencing have been utilized to provide detailed profiles of the structure and diversity of the microbiota residing on the mucosal surface of fish (Ghanbari *et al.*, 2015).

The gut microbiome composition has also become an important biomarker for understanding the influence of stress in fish (Llewellyn *et al.*, 2014), as numerous stressful stimuli have been shown to alter the microbiome composition (Xia *et al.*, 2014; Gaulke *et al.*, 2016). The gut microbiome composition can provide insights into the ecology and physiology of fish in a range of areas such as ecological speciation (Sevellec *et al.*, 2014), the biology of migratory fish (Llewellyn *et al.*, 2016), trophic interactions within ecosystems (Ingerslev *et al.*, 2014) and adaptation to extreme environments (Song *et al.*, 2016).

There are a number of challenges currently facing fish microbiome research. At present, the majority of data regarding the microbiome composition in wild teleost fish originates from laboratory models (Tarnecki *et al.*, 2017). More studies are required to see if captive-reared animals provide a reliable analogue for wild populations. Standardized protocols for collecting and generating microbiome data are also lacking, which could restrict progress as several processes have the potential to introduce differential bias in microbiota profiles (Salipante *et al.*, 2014; Hart *et al.*, 2015). Adopting a framework of robust, quality-controlled protocols (*e.g.* similar to human microbiome research; Methé *et al.*, 2012) would be of great benefit. In addition, there is currently a lack of non-invasive protocols for conducting longitudinal or repeated sampling of the gut microbial community in individual fish over time. The application of rectal swabs (Budding *et al.*, 2014) for sampling the vent of

fish could provide a non-invasive strategy for collecting microbiome data from individuals over time. Finally, time-series data could also enhance our knowledge in terms of the functional aspects of host lifecycles and the stability and resilience of microbiota (Goodrich *et al.*, 2014).

## SURVEY TOOLS

### FIELD-BASED SURVEYS

Fish population assessments are conducted using a wide range of techniques; the advantages, limitations, personnel requirements and health and safety considerations of each are presented in Table 1. It is encouraging to note that even well-established methods such as hydroacoustics are continually being improved, while emerging tools such as eDNA are beginning to be included in routine monitoring. We suggest that integrating methods and data series are key priorities for future research in this field.

In large and complex habitats it is often the case that a suite of survey methodologies has to be employed to sample different times, habitats and species effectively. Indeed, an advantage of field-based surveys is the ability to generate information from both fishery-independent (Nash *et al.*, 2016) and fishery-dependent (Shin *et al.*, 2010) data. The availability of a diversity of methodologies, however, can make the task of assessment in these habitats even more costly; issues also remain over how to use often disparate data types to develop a sound understanding of a fishery. Integrating methods represents a key means of improving data resolution from such field surveys. For instance, methods such as eDNA and hydroacoustics sampling provide comparatively fast and non-invasive estimates of fish community structure and biomass. To obtain a thorough understanding of fish populations,

however, this information must be combined with fish age, size and health data obtained *via* destructive sampling (*e.g.* gill netting). As yet, there are no structured, universally agreed guidelines on which methods should be integrated to obtain a thorough assessment of population dynamics from a specific habitat type.

Fish-survey methodologies are typically determined at a national level, making international comparisons of data extremely challenging. In recent years, standardized protocols initiated through the European Union Water Framework Directive (E.U., 2000) have facilitated Europe-wide assessments of fish community structure. Such international standardization is essential when assessing the anthropogenic effects on fish (Gordon *et al.*, 2018) and we recommend that efforts are made to make national datasets available using standardized metadata and biodiversity information, ideally via open sharing platforms (*e.g.* [www.freshwaterplatform.eu](http://www.freshwaterplatform.eu)).

## HISTORICAL RECORDS

Historical records (*e.g.* catch records) can also be useful in helping to extrapolate population data back into the recent past. Libraries and historical societies often hold picture archives and these images can in some instances be used as a form of historical survey data to provide information on past community composition and size distributions (McClenachan, 2009). Historical records of catch data are typically held by government agencies or can be found in local archives (*e.g.* angling club logs) and corporate records. Such data have been used successfully to reconstruct fish populations back to the late 1800s (Thurstan & Roberts, 2010; Thurstan *et al.*, 2010). Catch reconstruction approaches can also provide useful insights into fishery trends that may not be apparent from data reported only by the U.N. Food and Agriculture Organization (FAO; Smith & Zeller, 2015; Zeller *et al.*, 2015). Although limited

to the information that is still available and subject to the often-unidentifiable biases of the individuals who originally recorded the data, such data can provide a unique way to extrapolate population data back in time.

## STATISTICAL AND MODELLING TOOLS

### BAYESIAN METHODS

Reliable estimates of demographic variables (*e.g.* abundance, survival, growth rates and fecundity) and an understanding of the processes that regulate these variables are fundamental for sustainable management of fish populations. To understand the ecological processes to truly inform policy, however, researchers must use multiple data sources, provide links between management actions and population responses and also estimate uncertainty as a prerequisite to making forecasts that provide useful information. Bayesian methods in ecology and conservation biology are now increasingly being used to explore these links, for example, in stable-isotope analyses. Indeed, the Bayesian framework provides an intuitive method for estimating parameters, expressing uncertainty in these estimates and allows for the incorporation of as much or as little existing data or prior knowledge that is available (Ellison, 2004). To develop the use of this specific framework in fish ecology and management, however, there is a need to educate and train fish biologists in the use of Bayesian principles and methods.

### INDIVIDUAL-BASED MODELS

Individual-based models (IBM) are process-based mechanistic computer models that simulate emergent properties of fish biology, behaviour, traits or group characteristics, based on simple heuristic functions and their use has grown exponentially ( DeAngelis & Mooij, 2005) as computational power has increased (DeAngelis & Grimm, 2014). Several separate individual-based models were presented at the 50th Anniversary Symposium of The Fisheries Society of the British Isles and with continued increases in computational power, IBMs look set to offer powerful new avenues for population research (DeAngelis & Grimm, 2014) in computationally challenging multifactor systems such as fish ecotoxicology ( Mintram *et al.*, 2017). Additionally, a variety of tools now exist which provide for the easier creation of new models, such as various R packages ([www.r-project.org](http://www.r-project.org)) and programmable environments (*e.g.* NetLogo; [www.ccl.northwestern.edu/netlogo](http://www.ccl.northwestern.edu/netlogo)). Programmes such as R, however, are sometimes not intuitive to new users and so additional training for fisheries scientists and collaborations between scientists from different computational and statistical backgrounds would be advantageous. For more robust future application of IBMs within fisheries science, there is a need for more assessment of the relative strengths and weaknesses (and potential availability and future development) of the different models.

Integration with environmental data is a pertinent issue when modelling and is becoming easier through developments in geographic information systems (GIS) and other programming environments (such as R), which now include procedures and libraries for use in ecological work. One example is the use of food-web models that integrate environmental data ( Christensen & Walters, 2004) and coral-reef ecosystem modelling methods ( Rogers *et al.*, 2014; Weijerman *et al.*, 2015). A hindrance to the integration of environmental data into fisheries science is that it can be difficult to find and access data sources, although availability and accessibility of such data is improving ([www.worldclim.org](http://www.worldclim.org)). The existence of a central node or hub with paths to these data sources would be useful.

377

378

## TISSUE ANALYSIS

379

### 380 STABLE-ISOTOPE ECOLOGY

381

382       Stable isotopes are now routinely used to quantify the trophic ecology (Boecklen *et*  
383 *al.*, 2011) and migration history (Trueman *et al.*, 2012) of fish, or to identify community level  
384 patterns in food-web structure and resource use (Layman *et al.*, 2012). Although the  
385 technique is still in its relative infancy, stable-isotope ecology (or stable-isotopes analysis,  
386 SIA) has advanced much in recent decades. Below we outline four areas of rapid  
387 development with potential to enhance the applicability of this tool to studies of fish biology.

388

### 389 BIOCHEMICAL MECHANISM

390

391       The relationship between the isotopic composition of a consumer's tissues and that of  
392 its prey is fundamental to all applications of stable isotopes in ecology. While general  
393 principles are clear [*i.e.* faster reaction rates and preferential incorporation of light isotopes  
394 into excretory metabolites a process termed trophic fractionation (DeNiro & Epstein, 1977)],  
395 however, the precise mechanisms leading to fractionation and, particularly, the extent of  
396 isotopic fractionation expected under differing physiological conditions cannot currently be  
397 predicted, primarily due to the complexity of amino-acid biochemistry. Uncertainties  
398 associated with the isotopic expression of tissue composition and relative rates of tissue  
399 growth and regeneration further complicate the interpretation of stable isotope values in  
400 ecology. Recent information gained from compound-specific isotope analysis (*i.e.* assessing

isotopic compositions of single amino acids), however, is beginning to shed light on the fractionation process (McMahon & McCarthy, 2016).

#### POPULATION-LEVEL DATA

The distribution of isotopic compositions of individuals within a population (often termed the isotopic niche; Newsome *et al.*, 2007) has been proposed as a powerful comparative measure of population-level ecological characters. In addition to individual variability in consumers, however, the distribution of isotopic compositions in a population is influenced by spatial and temporal variations in the isotopic composition of primary production, temporal variability within trophic linkages and differential rates of growth and isotopic assimilation (Gorokhova, 2017). Very few studies have attempted to combine ecological and food web theory with isotope systematics to explore the sensitivity of community isotopic metrics to changes in food web structure and function.

#### ISOBANK

To date, applications of stable isotopes to fish biology have predominantly focussed on analyses of specific populations or communities. The absence of a centralized, open-access repository for stable-isotope data restricts the opportunity for syntheses or meta-analyses of stable-isotope data (Pauli *et al.*, 2017). Recent efforts to address this have found broad support from the stable isotope research community (Pauli *et al.*, 2017) and would be especially beneficial to fish biologists due to the large amount of fish-isotope data currently available. Defining an ontology of stable-isotope metadata, information required to describe and interpret isotope data, for fish biologists is an immediate requirement in this regard.

## MARINE ISOSCAPES

The stable-isotope ratios of a consumer's tissue encode the resources (water, air, prey *etc.*) it was using when that tissue was formed. As such, provided one has access to a suite of isotopic baseline measurements (*e.g.* water, plants and primary consumers), it is possible to trace an organisms route through space and time up to the point of capture (Trueman *et al.*, 2012). Creation of a practically useful isoscape requires relatively dense sampling of a reference organism across space (and potentially time). Bulk stable-isotope analyses are now routine, commonly available globally and relatively cheap and regional marine-isoscape models are being developed at a rapid rate (MacKenzie *et al.*, 2014; Kurle & McWhorter, 2017). In the open ocean, sample-based isoscapes are difficult to develop, but progress is being made in isotope-enabled global biogeochemical models (Magozzi *et al.*, 2017), offering temporal and spatial models of expected isotopic variability at global scales. Improving the precision, accuracy and availability of these baseline measurements will increase the robustness and precision of isotope-based estimates animal position.

## ARCHAEOLOGICAL MATERIAL

Archaeological material can allow an otherwise impossible snapshot into past populations. Traditional morphological approaches can provide age distributions and species ranges, and, with the rapid development of biomolecular archaeology in the past 20 years, many of the techniques used to explore modern fish populations can now be used to look into the past. From ancient DNA to proteomics and isotopes to lipids, a wide range of biomolecules have been recovered and explored from archaeological material (Orton, 2016). For example, compound-specific isotope analysis has the potential to track trophic level



changes through time (McClelland & Montoya, 2002; Naito *et al.*, 2016). Population genetics of extinct populations have been successfully explored in terrestrial animals (Chang & Shapiro, 2016; Murray *et al.*, 2017) and these same techniques can be used on fish bones to reconstruct past genetics (Iwamoto *et al.*, 2012; Ólafsdóttir *et al.*, 2014). Ideally these data will be used to understand environmental and anthropogenic effects on fish populations and how modern fish populations might respond to climate change and fishing pressures.

A major barrier to the use of archaeological fish material is the fact that less than 10% of fish bones are identified to species (Wheeler & Jones, 1989; Gobalet, 2001) and much of what is identified is buried in the grey literature of archaeological reports that are often printed in small quantities and not digitized (Linden & Webley, 2012). This makes the material relevant to an ecological question very difficult to find. Archaeologists are working towards ways to improve the amount of bones identified by better reference collections and education on fish bones (National Zooarchaeological Reference Resource, Nottingham's Archaeological Fish Resource; Vertebra@UWF) and on creating searchable databases of archaeological material (Callou, 2009; Kansa, 2010). In addition, new ZooMS (Zooarchaeology by mass spectrometry) techniques are being explored to quickly identify even small bones and scales to species using peptide mass fingerprinting (Richter *et al.*, 2011) which will allow even more material to be identified in a useful way for those working on understanding fish populations. In the near future, it should be possible for modern fish biologists, in conjunction with archaeologists, to ask direct questions of past populations (Van Neer & Ervynck, 2010).

## GENERAL TOPICS IDENTIFIED AS APPLICABLE ACROSS ALL THEMES

MANAGEMENT OF DATA: INTEGRATION, CALIBRATION AND STANDARDIZATION

Progression of an integrated management framework for data classification, characterization, storage and accessibility would be a valuable resource for fish and fisheries biologists. FishBase, which at the time of writing contains information regarding 33 600 fishes, involving 2290 collaborators and receives over 600 000 visits per month, is an example of the potential for such a resource ([www.fishbase.org](http://www.fishbase.org); Froese & Pauly, 2017). A single database for all types of fish data (for example, DNA, tagging, isotopes, diet) is probably unworkable, but the advent of application programming interfaces (API) and analytical software which allows automated querying across multiple databases represents an unprecedented opportunity to access a wealth of global data. Indeed, we suggest that more data (such as those discussed here) could be integrated into FishBase. Such resources, however, require significant funding and long-term commitment from governments and trans-national organisations, *e.g.* the North Atlantic Salmon Conservation Organisation (NASCO).

## PUBLIC ENGAGEMENT, EDUCATION AND OUTREACH

Scientific engagement with the public is essential to effect meaningful societal change or to ensure a wider consensus is made around new discoveries or ethical considerations. Additionally, however, the power of the public as a tool in science is also being increasingly recognized. Crowdfunding, whereby a scientist requests small amounts of money from a large number of interested individuals to successfully launch a project, potentially provides a powerful new way to raise funds, overcoming some of the difficulties of raising money from traditional grant bodies, especially for early career researchers or those in developing countries (Wheat *et al.*, 2013).

In addition to funding science, the public can also actively engage in the process of research directly through citizen science projects. Whilst research conducted by non-professionals is certainly not a new concept, the numbers of projects involving citizen scientists are growing, especially in the fields of environmental science and ecology (Silverton, 2009). Through catch records of amateur anglers and commercial net-fishery data extending back many years, research into fish and fisheries is uniquely placed to benefit from citizen science projects (Stuart-Smith *et al.*, 2013), which have effectively spanned generations of contributors. Similarly, REEF ([www.reef.org](http://www.reef.org)) has been collecting reef-fish diversity and abundance data from trained volunteer divers for 27 years and the data have been successfully leveraged in hundreds of publications (Stallings, 2009; Serafy *et al.*, 2015). Citizen science can also help achieve important social outcomes, *e.g.* in establishing sustainable fisheries and marine protected areas (MPA) (Bonney *et al.*, 2014). As with crowdfunding, the best examples of citizen science typically encourage deeper engagement with the public and offer a pathway to the democratisation of science.

## FISHERIES POLICY AND GOVERNANCE

Conserving critical habitats is central to the sustainable management of fish species and populations. MPAs, networks of MPAs and marine conservation zones (MCZ) are widely accepted management tools for fish and other marine organisms that have been established in many countries (Harborne *et al.*, 2008; OSPAR, 2013). The design of MPA networks could, however, benefit greatly from the integration of traditional survey data, along with modelling and connectivity data (Botsford *et al.*, 2009; Grüss *et al.*, 2014). From a social science perspective, there is a need to better understand public perceptions of marine-related conservation issues, *e.g.* fishery regulations, MPAs and MCZs and to incorporate these data

into fisheries policy and governance frameworks. For example, there is high public support for MPAs, with surveys showing that people desire around 40% of the UK's marine waters to be protected (Hawkins *et al.*, 2016). But, while the public appears to realize that in reality levels of coverage are well below 40%, there is still a substantial disconnect between perceived coverage of highly protected UK MPAs (11%) and actual MPA coverage (<0.1%); ultimately, this means that people believe the UK oceans receive a higher level of conservation than in reality they do (Hawkins *et al.*, 2016). Developing and implementing effective policies for fisheries management remains challenging because of the complexities of fisheries and the socio-political landscape under which they typically operate (Jentoft & Chuenpagdee, 2009). The establishment of guidelines or frameworks for fisheries policy and governance (FAO, 2015), however, have the potential to better address these challenges and provide appropriate implementable solutions.

## CONCLUSIONS

Across all five of the research themes identified here, it is clear that innovative and novel tools are being employed to understand all aspects of the biology of fish populations. Notwithstanding, the authors call for the continued development of these new and emerging techniques. In particular, there is a need for better integration of these methods and resulting data, to inform scientifically sound management and conservation of fish populations. It should be noted, however, that not infrequently, revolutionary methods have been promoted as providing the ability to offer unprecedented novel answers to long-standing practical problems. Unfortunately, the danger is that such methods can (by their novelty and the excitement surrounding them), blinker scientists into posing questions that showcase the methodology, rather than the biology [*e.g.* the plethora of papers that emerged in the early

1990s extolling the virtues of the random amplified polymorphic DNA (RAPD) technique]. The potentially reduced power of using any technique on its own (new or otherwise), in isolation of other apparently antiquated methods can turn out to be unnecessarily restrictive. Every technique has its limitations, but often the restrictions of one tool can be substantially alleviated by the inclusion of another approach ( Goodwin *et al.*, 2016; Nielson *et al.*, 2017), the marriage of which can provide a new angle for researching challenging biological problems. It is important that both traditional and emerging tools remain in the toolbox of fish biology research.

Likewise, when genetic-based assignment became popular, many researchers naively believed the days of tagging fish were over. It is now realized that due to the many stochastic drivers of population structure, genetic stock identification-based methodologies such as genetic assignment, do not always succeed. In such cases, there remains a significant role for tagging in fish and fisheries research. As tag sizes decrease and the deleterious effects of tag insertions on fish also decrease, we can anticipate that genetics and tagging will both continue to have a role to play. The importance of the relative roles of each technique will depend on the questions being addressed, the population structure of the study species and the scale of the questions being assessed.

A final example, which highlights the importance of applying inter-disciplinary and complimentary tools for understanding fish populations, was a five-year, multi-agency, E.U. funded project investigating the migration and distribution of *S. salar* in the north-east Atlantic (the SALSEA project; NASCO 2008). The purpose was to understand not just where *S. salar* go, but what they eat, migration routes to feeding grounds and which waters and regions they pass through. The SALSEA project used a combination of genetics (microsatellites), SIA, at-sea trawls, tagging and gut contents analysis to assess the movements and diet of this species across the north-east Atlantic Ocean. As a result of

applying these combined approaches, *S. salar* post-smolt movements have been confidently ascertained (Gilbey *et al.*, 2017). Nonetheless, even while this comprehensive study was being finalized, a similarly broad-ranging study was also being undertaken using SNPs (Bourret *et al.*, 2013). Arguably, this method offers both the potential for finer levels of stock discrimination and the ability to better explore patterns among functional loci, which may make microsatellite-based analysis redundant within a short period of time (though see Narum *et al.*, 2008).

Thus, the authors consider the continued development of emerging tools, together with the use of multiple methodologies and inter-disciplinary approaches, to represent the best avenues for further improving our understanding of fish populations. We implore scientists from unrelated fields to collaborate on such projects. The 50th Anniversary Symposium of The Fisheries Society of the British Isles represented one such event, where fish-focused researchers across diverse fields, came together to advance the state of fish biology.

We thank the organising committee of the 50th Anniversary Symposium of The Fisheries Society of the British Isles for enabling the working group discussion that led to the development of this review. Thanks also to the University of Exeter for hosting the symposium and to the numerous sponsors for funding its success. J.R.S. acknowledges the support of the SAMARCH project, an E.U. funded France Channel–Manche England Interreg initiative.

## References

- Allendorf, F. W., Hohenlohe, P. A. & Luikart, G. (2010). Genomics and the future of conservation genetics. *Nature Reviews Genetics*, **11**, 697–709. doi:10.1038/nrg2844
- Amann, R. I., Ludwig, W. & Schleifer, K. H. (1995). Phylogenetic identification and *in situ* detection of individual microbial cells without cultivation. *Microbiological Reviews*, **59** 143-169.
- Austin, B. (2006). The bacterial microflora of fish, revised. *The Scientific World Journal* **6**, 931-945. doi: 10.1100/tsw.2006.181
- Austin, C. B., Hua Tan, M., Harrison, K. A., Peng Lee, Y., Croft, L. J., Sunnucks, P., Pavlova, A. & Ming Gan, H. (2017). *De novo* genome assembly and annotation of Australia's largest freshwater fish, the Murray cod (*Maccullochella peelii*) from Illumina and Nanopore sequencing reads. *Gigascience* gix063. doi: 10.1093/gigascience/gix063
- Benestan, L., Moore J.-S., Sutherland, B. J. G., Le Luyer, J., Maaroufi, H., Rougeux, C., Normandeau, E., RYcroft, N., Atema, J., Harris, L. N., Tallman, R. F., Greenwood, S. J., Clark, F. K. & Bernatchez, L. (2017). Sex matters in massive parallel sequencing: Evidence for biases in genetic parameter estimation and investigation of sex determination systems. *Molecular Ecology* (online). doi.org/10.1111/mec.14217
- Blanchet, S., Prunier, J. G. & De Kort, H. (2017). Time to go bigger: emerging patterns in macrogenetics. *Trends in Genetics* **9**, 579–580. doi: 10.1016/j.tig.2017.06.007
- Boecklen, W. J., Yarnes, C. T., Cook, B. A., & James, A. C. (2011). On the use of stable isotopes in trophic ecology. *Annual Review of Ecology, Evolution and Systematics*, **42**, 411–440. doi.org/10.1146/annurev-ecolsys-102209-144726

- Bograd, S. J., Block, B. A., Costa, D. P. & Godley, B. J. (2010). Biologging technologies: new tools for conservation. Introduction. *Endangered Species Research* **10**, 1–7. doi: 10.3354/esr00269
- Bonney, R., Shirk, J. L., Phillips, T. B., Wiggins, A., Ballard, H. L., Miller-Rushing, A. J. & Parrish, J. K. (2014). Next steps for citizen science. *Science* **343**, 1436–1437. doi: 10.1126/science.1251554
- Botsford, L. W., Brumbaugh, D. R., Grimes, C., Kellner, J. B. Largier, J., O'Farell, M. R., Ralston, S., Soulanile, E. & Wespestad, V. (2009). Connectivity, sustainability and yield: bridging the gap between conventional fisheries management and marine protected areas. *Reviews in Fish Biology Fisheries* **19**, 69–95. doi: 10.1007/s11160-008-9092-z
- Bourret, V., Kent, M. P., Primmer, C. R., Vasermägi, A., Karlsson, S., Hindar, K., McGinnity, P., Verspoor, E., Bernatches, L. & Lien, S. (2013). SNP-array reveals genome-wide patterns of geographical and potential adaptive divergence across the natural range of Atlantic salmon (*Salmo salar*). *Molecular Ecology*, **22**, 532–551.
- Budding, A. E., Grasman, M. E., Eck, A., Bogaards, J. A., Vandenbroucke-Grauls, C. M., van Bodegraven, A. A. & Savelkoul, P. H. (2014). Rectal swabs for analysis of the intestinal microbiota. *PLoS One* **9**, e101344. doi.org/10.1371/journal.pone.0101344
- Chang, D. and Shapiro, B. (2016). Using ancient DNA and coalescent-based methods to infer extinction. *Biology Letters*, **12**, 20150822.
- Chistiakov, D. A., Hellemans, B. & Volckaert, F.A.M. (2006). Microsatellites and their genomic distribution, evolution, function and applications: a review with special reference to fish genetics. *Aquaculture*, **255**, 1–29.
- Christensen, V. & Walters, C.J. (2004) Ecopath with Ecosim: methods, capabilities and limitations. *Ecological Modelling*, **172**, 109-139.



- Clusa, L., Ardura, A., Fernández, S., Roca, A. A. & García-Vázquez, E. (2017). An extremely sensitive nested PCR-RFLP mitochondrial marker for detection and identification of salmonids in eDNA from water samples. *PeerJ* **5**, p.e3045.
- Cooke, S. J., Brownscombe, J. W., Raby, G. D., Broell, F., Hinch, S. G., Clark, T. D., & Semmens, J. M. (2016). Remote bioenergetics measurements in wild fish: opportunities and challenges. *Comparative Biochemistry and Physiology A* **202**, 23-37.
- Cooke, S. J., Midwood, J. D., Thiem, J. D., Klimley, P., Lucas, M. C., Thorstad, E. B., Eiler, J., Holbrook, C. & Ebner, B. C. (2013). Tracking animals in freshwater with electronic tags: past, present and future. *Animal Biotelemetry*, **1**, 5.
- Dahlgren, C. P., Buch, K., Rechisky E. & Hixon M. A. (2016). Multiyear tracking of Nassau grouper spawning migrations. *Marine and Coastal Fisheries: Dynamics, Management and Ecosystem Science*, **8**, 522–535. DOI: 10.1080/19425120.2016.1223233.
- Davey, J. W. & Blaxter, M. L. (2011). RADSeq: next-generation population genetics. *Briefings in Functional Genomics* **9**, 416–423.
- DeAngelis D. L. & Grimm, V. (2014). Individual-based models in ecology after four decades. *F1000Prime Reports*, **6**, 39. doi:10.12703/P6-39
- DeAngelis, D. L. & Mooij, W. M. (2005). Individual-based modelling of ecological and evolutionary processes. 1. *Annual Review of Ecology, Evolution and Systematics*, **36**, 147–168.
- DeNiro, M. J. & Epstein, S. (1977). Mechanism of carbon isotope fractionation associated with lipid synthesis. *Science*, **197**, 261–263.

- Doi, H., Uchii, K., Takahara, T., Matsushashi, S., Yamanaka, H. & Minamoto, T., (2015). Use of droplet digital PCR for estimation of fish abundance and biomass in environmental DNA surveys. *PLoS One* **10**, e0122763.
- Ekblom, R. & Wolf, J. B. W. (2014). A field guide to whole-genome sequencing, assembly and annotation. *Evolutionary Applications*, **7**, 1026–1042.
- Ellegren, H. & Galtier, N. (2016). Determinants of genetic diversity. *Nature Reviews Genetics*, **17**, 422–433. doi.org/10.1038/nrg.2016.58
- Ellis, J. S., Gilbey, J., Armstrong, A., Balstad, T., Cauwelier, E., Cherbonnel, C., Consuegra, S. et al. (2011). Microsatellite standardization and evaluation of genotyping error in a large multi-partner research programme for conservation of Atlantic salmon (*Salmo salar* L.). *Genetica*, **139**, 353–367.
- Ellison, A. M. (2004). Bayesian Inference in Ecology. *Ecology Letters*, **7**, 509–520.
- E.C. (2000). Directive 2000/60/EC, Establishing a framework for community action in the field of water policy. *Official Journal of the European Communities*, **327**, 1–71. Available at [www.lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2000:327:0001:0072:EN:PDF](http://www.lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2000:327:0001:0072:EN:PDF)
- FAO. (2014). *The State of World Fisheries and Aquaculture Opportunities and Challenges* 2014. Rome: FAO. Available at <http://www.fao.org/3/a-i3720e.pdf>
- FAO. (2015). *Voluntary Guidelines for Securing Sustainable Small-Scale Fisheries in the Context of Food Security and Poverty Eradication*, Rome: FAO. Available at <http://www.fao.org/fishery/ssf/guidelines/en>
- Ferguson, M. M. & Danzmann, R. G. (1998). Role of genetic markers in fisheries and aquaculture: useful tools or stamp collecting? *Canadian Journal of Fisheries and Aquatic Sciences*, **55**, 1553–1563.
- Freeland, J. R. (2016). The importance of molecular markers and primer design when characterizing biodiversity from environmental DNA. *Genome*, **60**, 358–374.

- Gaulke, C. A., Barton, C. L., Proffitt, S., Tanguay, R. L. & Sharpton, T. J. (2016). Triclosan exposure is associated with rapid restructuring of the microbiome in adult zebrafish. *PloS One*, **11**. e.0154632
- Ghanbari, M., Kneifel, W. & Domig, K. J. (2015). A new view of the fish gut microbiome: advances from next-generation sequencing. *Aquaculture*, **448**, 464-475. doi: 10.1016/j.aquaculture.2015.06.033
- Gilbey J., Coughlan, J., Wennevik, V., Prodöhl, P. Stevens, J. R., Garcia de Leaniz, C., Ensing, D., Cauwelier, E., Cherbonnel, C., Consuegra, S., Coulson, M. W., Cross, T. F., Crozier, W., Dillane, E., Ellis, J. S., García-Vázquez, E., Griffiths, A. M., Gudjonsson, S., Hindar, K., Karlsson, S., Knox, D., Machado-Schiaffino, G., Meldrup, D., Nielsen, E. E., Ólafsson, K., Primmer, C. R., Prusov, S., Stradmeyer, L., Vähä, J.-P., Veselov, A. J., Webster, L. M. I., McGinnity, P. & Verspoor, E. (2017). A microsatellite baseline for genetic stock identification of European Atlantic salmon (*Salmo salar* L.). *ICES journal of Marine Science* (online). doi.org/10.1093/icesjms/fsx184
- Gobalet, K.W. (2001). A critique of faunal analysis; inconsistency among experts in blind tests. *Journal of Archaeological Science*, **28**, 377e386.
- Goodrich, J. K., Di Rienzi, S. C., Poole, A. C., Koren, O., Walters, W. A., Caporaso, J. G., Knight, R. & Ley, R. E. (2014). Conducting a microbiome study. *Cell* **158**, 250–262. doi: 10.1016/j.cell.2014.06.037
- Goodwin, J. C. A., King, R. A., Jones, J. I., Ibbotson, A. & Stevens, J. R. (2016). A small number of anadromous females drive reproduction in a brown trout (*Salmo trutta*) population in an English chalk stream. *Freshwater Biology*, **61**, 1075–1089.
- Gordon, T. A. C., Harding, H. R., Clever, F. K., Davidson, I. K., Davison, W., Montgomery, D. W., Nedelec, S. L., Weatherhead, R. C., Windsor, F. M., Armstrong, J. D.,

- Bardonnet, A., Bergman, E., Britton, J. R., Côté, I. M., D'Agostino, D., Greenberg, L. A., Harborne, A. R., Kahilainen, K. K., Metcalfe, N. B., Mills, S. C., Milner, N. J., Mittermayer, F. H., Montorio, L., Prokkola, L. A., Rutterford, L. A., Salvanes, A. G. V., Simpson, S. D., Vainikka, A., Pinnegar, J. K., Santos, E. M. (2018). Fishes in a changing world: learning from the past to promote sustainability of fish populations. *Journal of Fish Biology*, (In review, this issue).
- Gorokhova E. (2017). Individual growth as a non-dietary determinant of the isotopic niche metrics. *Methods in Ecology and Evolution* (online). doi.org/10.1111/2041-210X.12887
- Gruss, A., Robinson, J., Heppell, S. S., Heppell, S. A., & Semmens, B. X. (2014). Conservation and fisheries effects of spawning aggregation marine protected areas: what we know, where we should go and what we need to get there. *ICES Journal of Marine Science*, **71**, 1515–1534. doi.org/10.1093/icesjms/fsu038
- Guillot, G., Leblois, R., Coulon, A. & Frantz, A. C. (2009). Statistical methods in spatial genetics. *Molecular Ecology* **18**, 4734–4756. doi:10.1111/j.1365-294X.2009.04410.x
- Halfyard, E. A., Webber, D., Del Papa, J., Leadley, T., Kessel, S. T., Colborne, S. F. and Fisk, A. T. (2017), Evaluation of an acoustic telemetry transmitter designed to identify predation events. *Methods in Ecology and Evolution* **8**, 1063–1071. doi:10.1111/2041-210X.12726.
- Hänfling, B., Lawson Handley, L., Read, D. S., Hahn, C., Li, J., Nichols, P., Blackman, R. C., Oliver, A. & Winfield, I. J., 2016. Environmental DNA metabarcoding of lake fish communities reflects long-term data from established survey methods. *Molecular Ecology* **25**, 3101-3119.

- Harborne, A. R., Mumby, P. J., Cappel, C. V., Dahlgren, C. P., Micheli, F., Holmes, K. E., Sanchirico, J. E., Broad, K., Elliot, I. A., Brumbaugh, D. R. (2008) Reserve effects and natural variation in coral reef communities. *Journal of Applied Ecology* **45**, 1010–1018.
- Hart, M. L., Meyer, A., Johnson, P. J. & Ericsson, A. C. (2015). Comparative evaluation of DNA extraction methods from feces of multiple host species for downstream next-generation sequencing. *PloS ONE*, **10**, e0143334. doi: 10.1371/journal.pone.0143334
- Hawkins, J. P., O’Leary, B. C., Bassett, N., Peters, H., Rakowski, S., Reeve, G. & Roberts, C. M. (2016). Public awareness and attitudes towards marine protection in the United Kingdom. *Marine Pollution Bulletin*, **111**, 231–236.
- Hayden, E. C. (2015). Pint-sized DNA sequencer impresses first users. *Nature*, **521**, 15-16.
- Hays, G. C., Bradshaw, C. J. A., James, M. C., Lovell, P. & Sims, D. W. (2007). Why do Argos satellite tags deployed on marine animals stop transmitting? *Journal of Experimental Marine Biology and Ecology*, **349**, 52–60.
- Helyar, S. J., Hemmer-Hansen, J., Bekkevold, D., Taylor, M. I., Ogden, R., Limborg, M. T., Cariani, A., Maes, G. E., Dopere, E., Carvalho, G. & Nielsen, E. E. (2011). Application of SNPs for population genetics of nonmodel organisms: new opportunities and challenges: analytical approaches. *Molecular Ecology Resources* **11**, 123–136.
- Heupel, M. R., Semmens, J. M. & Hobday, A. J. (2006) Automated acoustic tracking of aquatic animals: scales, design and deployment of listening station arrays. *Marine and Freshwater Research* **57**, 1–13.
- Hight, B. V. & Lowe, C. G. (2007). Elevated body temperatures of adult female leopard sharks, *Triakis semifasciata*, while aggregating in shallow nearshore embayments:

evidence for behavioral thermoregulation? *Journal of Experimental Marine Biology and Ecology* **352**, 114–128.

Hussey, N. E., Kessel, S. T., Aarestrup, K., Cooke, S. J., Cowley, P. D., Fisk, A. T., Harcourt, R. G., Holland, K. N., Iverson, S. J., Kocik, J. F., Flemming, J. E. M. & Whoriskey, F. G. (2015). Aquatic animal telemetry: a panoramic window into the underwater world. *Science* **348**, 1221–1255642. doi: 10.1126/science.1255642

Ingerslev, H. C., von Gersdorff Jørgensen, L., Strube, M. L., Larsen, N., Dalsgaard, I., Boye, M. & Madsen, L. (2014). The development of the gut microbiota in rainbow trout (*Oncorhynchus mykiss*) is affected by first feeding and diet type. *Aquaculture*, **424**, 24–34. doi: 10.1016/j.aquaculture.2013.12.032

Iwamoto, E. M., Myers, J. M. & Gustafson, R. G. (2012). Resurrecting an extinct salmon evolutionary significant unit: archived scales, historical DNA and implication for restoration. *Molecular Ecology*, **21**, 1567-1582.

Jansen, H. J., M., Liem, S. A., Jong-Raadsen, S., Dufour, F.-A., Weltzien, W., Swinkels, A., Koelewijn, A. P., Palstra, B., Pelster, H. P., Spaik, G. E., Van den Thillart, Dirks, R. P. & Henkel, C. V. (2017). Rapid de novo assembly of the European eel genome from nanopore sequencing reads. *bioRxiv* 101907. doi: 10.1101/101907

Jentoft, S., & Chuenpagdee, R. (2009). Fisheries and coastal governance as a wicked problem. *Marine Policy*, **33**, 553–560. doi: 10.1016/j.marpol.2008.12.002

Jepsen, N., Thorstad, E. B., Havn, T. & Lucas, M. C. (2015). The use of external electronic tags on fish: an evaluation of tag retention and tagging effects. *Animal Biotelemetry*, **3**, 49. 10.1186/s40317-015-0086-z.

Kansa, E. C. (2010). Open context in context: cyberinfrastructure and distributed approaches to publish and preserve archaeological data. *The SAA Archaeological Record*, **10**, 12–16.

798 Kurle, C. M. & McWhorter, J. (2017) Spatial and temporal variability within marine isoscaes:  
799 implications for interpreting stable isotope data from marine systems. *Marine*  
800 *Ecology Progress Series*, **568**, 31–45. doi 10.3354/meps12045

801 Lacoursière- Roussel, A., Rosabal, M. & Bernatchez, L. (2016). Estimating fish abundance  
802 and biomass from eDNA concentrations: variability among capture methods and  
803 environmental conditions. *Molecular Ecology Resources*, **16**, 1401–1414.

804 Lauridsen, R. B., Moore, A., Privitera, L. Gregory, S. D., Beaumont, W. R. C. & Kavanagh,  
805 A. J. (2017). Migration behaviour and loss rate of trout smolts in the transitional  
806 zone between freshwater and saltwater In: *Proceedings of the 2nd International Sea*  
807 *Trout Symposium Sea Trout: Science & Management*. (Harris, G., ed.) pp. 292–307.  
808 Leicester: Matador

809 Laval, G. & Excoffier, L. (2004). SIMCOAL 2.0: a program to simulate genomic diversity  
810 over large recombining regions in a subdivided population with a complex history.  
811 *Bioinformatics*, **20**, 2485–7. DOI: 10.1093/bioinformatics/bth264.

812 Layman, C. A., Araujo, M. S., Boucek, R., Hammerschlag-Peyer, C. M., Harrison, E., Jud, Z.  
813 R., Matich, P., Rosenblatt, A. E., Vaudo, J. J., Yeager, L. A., Post, D. M. &  
814 Bearhop, S. (2012), Applying stable isotopes to examine food-web structure: an  
815 overview of analytical tools. *Biological Reviews* **87**, 545–562. doi:10.1111/j.1469-  
816 185X.2011.00208.x

817 Lea, J. S. E., Humphries, N. E., von Brandis, R. G., Clarke C. R. & Sims, D. W. (2016).  
818 Acoustic telemetry and network analysis reveal the space use of multiple reef  
819 predators and enhance marine protected area design. *Proceedings of the Royal*  
820 *Society B* **283** (online). DOI: 10.1098/rspb.2016.0717

821 Lien, S., Koop, B. F., Sandve, S. R., Miller, J. R., Matthew, P., Leong, J. S., Minkley, D. R.,  
 822 Zimin, A., Grammes, F., Grove, H. et al. (2016) The Atlantic salmon genome  
 823 provides insights into rediploidization. *Nature* **533**, 200–205.

824 Linden, M. V. & Webley, L. (2012). Introduction: Development-led archaeology in north-  
 825 west Europe. In, *Development-led Archaeology in North-West Europe* (Bradley, R.,  
 826 Haselgrove, C., Linden, M. V. & Webley, L., eds.). Oxford: Oxbow Books.

827 Llewellyn, M. S., Boutin, S., Hoseinifar, S. H. & Derome, N. (2014). Teleost microbiomes:  
 828 the state of the art in their characterization, manipulation and importance in  
 829 aquaculture and fisheries. *Frontiers in Microbiology* **5**. doi:  
 830 10.3389/fmicb.2014.00207

831 Llewellyn, M. S., McGinnity, P., Dionne, M., Letourneau, J., Thonier, F., Carvalho, G. R.,  
 832 Creer, S. & Derome, N. (2016). The biogeography of the Atlantic salmon (*Salmo*  
 833 *salar*) gut microbiome. *The ISME Journal* **10**, 1280–1284. doi:  
 834 10.1038/ismej.2015.189

835 MacDonald, A. J. & Sarre, S. D. (2017). A framework for developing and validating  
 836 taxon- specific primers for specimen identification from environmental DNA.  
 837 *Molecular Ecology Resources* **17**, 708–720.

838 MacKenzie, K. M., Longmore, C., Preece, C., Lucas, C. H., Trueman, C. N. (2014). Testing  
 839 the long-term stability of marine isoscapes in shelf seas using jellyfish tissues.  
 840 *Biogeochemistry*, **121**, 441–454. doi: 10.1007/s10533-014-0011-1

841 Macqueen, D. J., Primmer, C. R., Houston, R. D., Nowak, B. F., Bernatchez, L., Bergseth,  
 842 S., Davidson, W. S., Gallardo-escárate, C., Goldammer, T., Guiguen, Y., •  
 843 Patricia Iturra, Kijas, J. W., Koop, B. F., Lien, S., Martin, S. A. M., McGinnity, P.,  
 844 Montecino, M., Naish, K. A., Nichols, K. M., Ólafsson, K., Omholt, S. W., Palti, Y.,  
 845 Plastow, G. S., Rexroad 3<sup>rd</sup>, C. E., Rise, M. L., Ritchie, R. J., Sandve, S. R., Schulte,



P. M., Tello, A., Vidal, R., Vik, J. O., Wargelius, A., Yáñez, J. m. & the FAASG  
onsortium (2017). Functional Annotation of All Salmonid Genomes (FAASG): an  
international initiative supporting future salmonid research, conservation and  
aquaculture. *BMC Genomics* **18**, 1–9.

Magozzi, S. Yool, A., Van der Zanden, H. B., Wunder, M. B., Trueman, C. N. (2017). Using  
ocean models to predict spatial and temporal variation in marine carbon isotopes.  
*Ecosphere* **8**, e01763. doi: 10.1002/ecs2.1763

McClelland, J. W. and Montoya, J. P. (2002). Trophic relationships and the nitrogen isotopic  
composition of amino acids in plankton. *Ecology* **83**, 2173–2180.

McClenachan, L. (2009). Documenting loss of large trophy fish from the Florida Keys with  
historical photographs. *Conservation Biology*, **23**, 636–643.

McMahon, K. W. & McCarthy, M. D. (2016). Embracing variability in amino acid  $\delta^{15}\text{N}$   
fractionation: mechanisms, implications and applications for trophic ecology.  
*Ecosphere* **7**, e01511

Methé, B. A., Nelson, K. E., Pop, M., Creasy, H. H., Giglio, M. G., Huttenhower, C., Gevers,  
D., Petrosino, J. F., Abubucker, S., Badger, J. H. & Chinwalla, A.T. (2012). A  
framework for human microbiome research. *Nature* **486**, 215–221. doi:  
10.1038/nature11209

Methling, C., Tudorache, C., Skov, P. V. & Steffensen, J. F. (2011). Pop up satellite tags  
impair swimming performance and energetics of the European eel (*Anguilla*  
*anguilla*). *Plos One* **6**, e20797. doi: 10.1371/journal.pone.0020797

Mintram, K. S., Brown, A. R., Maynard, S. K., Thorbek, P. & Tyler, C. R. (2017). Capturing  
ecology in modeling approaches applied to environmental risk assessment of  
endocrine active chemicals in fish. *Critical Reviews in Toxicology*, DOI:  
10.1080/10408444.2017.1367756

- Miya, M., Sato, Y., Fukunaga, T., Sado, T., Poulsen, J. Y., Sato, K., Minamoto, T., Yamamoto, S., Yamanaka, H., Araki, H. & Kondoh, M. (2015). MiFish, a set of universal PCR primers for metabarcoding environmental DNA from fishes: detection of more than 230 subtropical marine species. *Open Science* **2**, 150088.
- Murchie, K.J., Shultz, A.D., Stein, J.A., Cooke, S. J., Lewis, J., Franklin, J., Vincent, G., Brooks, E. J., Claussen, J. E. & Philipp, D. P.(2015). Defining adult bonefish (*Albula vulpes*) movement corridors around Grand Bahama in the Bahamian Archipelago *Environmental Biology of Fishes* **98**, 2203–2212.  
doi.org/10.1007/s10641-015-0422-4
- Murray, G. G. R., Soares, A. E. R., Novak, B. J., Schaefer, N. K., Cahill, J. A., Baker, A. J., Shapiro, B. (2017). Natural selection shaped the rise and fall of passenger pigeon genomic diversity. *bioRxiv*. doi.org/10.1101/154294
- Musyl, M. K., Domeier, M. L., Nasby-Lucase, N., Brill, R. W., McNaughton, L. M., Swimmer, J. Y., Lutcavage, M. S., Wilson, S. G., Galuardi, B., Liddle, J. B. (2011). Performance of pop-up satellite archival tags. *Marine Ecology Progress Series*, **433**, 1–28.
- Naito, Y. I., Chikaraishi, Y., Drucker, D. G., Ohkouchi, N., Semal, P., Wißing, C. & Bocherens, H. (2016). Ecological niche of Neanderthals from Spy Cave revealed by nitrogen isotopes of individual amino acids in collagen. *Journal of Human Evolution* **93**, 82–90.
- Narum, S. R., Banks, M., Beacham, T. D., Bellinger, M. R., Campbell, M. R., DeKoning, J., Elz, A., Guthrie III, C. M., Kozfkay, C., Miller, K. M., Moran, P., Phillips, R., Seeb, L. W., Smith, C. T., Warheit, K., Young, S. F. & Garza, J. C. (2008). Differentiating salmon populations at broad and fine geographic scales with microsatellites and SNPs. *Molecular Ecology* **17**, 3464–3477.

896 Nash, K.L., Bijoux, J., Robinson, J., Wilson, S.K. & Graham, N.A.J. (2016) Harnessing  
897 fishery-independent indicators to aid management of data-poor fisheries: weighing  
898 habitat and fishing effects. *Ecosphere* **7**, e01362.

899 Nielsen, J. M., Clare, E. L., Hayden, B., Brett, M.T., Kratina, P. (2017). Diet tracing in  
900 ecology: method comparison and selection. *Methods in Ecology and Evolution*  
901 (online). doi.org/10.1111/2041-210X.12869

902 Neuenschwander, S. (2006) AQUASPLATCHE: a program to simulate genetic diversity in  
903 populations living in linear habitats. *Molecular Ecology Notes* **6** 583–585.

904 Newsome, S. D., del Rio, C. M., Bearhop, S. & Phillips D. L. (2007) A niche for isotopic  
905 ecology. *Frontiers in Ecology and the Environment* **5**, 429–436.

906 Ólafsdóttir, G. Á., Westfall, K. M., Edvardsson, R. & Pálsson, S. (2014). Historical DNA  
907 reveals the demographic history of Atlantic cod (*Gadus morhua*) in medieval and  
908 early modern Iceland. *Proceedings of the Royal Society B*, **281**, 20132976.

909 Orton, D. C. (2016). Archaeology as a tool for understanding past marine resource use and its  
910 impact. In *Perspectives on Oceans Past* (Schwerdtner Máñez, K. & Poulsen, B.,  
911 eds ), pp. 47–69. Dordrecht: Springer

912 Papastamatiou, Y. P., C. G. Meyer, F. Carvalho, J. J. Dale, M. R. Hutchinson & K. N.  
913 Holland. (2013). Telemetry and random-walk models reveal complex patterns of  
914 partial migration in a large marine predator. *Ecology* **94**, 2595–2606.

915 Paris, J. R., Stevens, J. R. & Catchen, J. M. (2017). Lost in parameter space: a road map for  
916 stacks. *Methods in Ecology and Evolution* **8**, 1360–1373. doi.org/10.1111/2041-  
917 210X.12775

918 Pauli, J. N., Newsome, S. D., Cook, J. A., Harrod, C., Steffan, S. A., Baker, C. J. O., Ben-  
919 David, M., Bloom, D., Bowen, G. J., Cerling, T. E. et al. (2017). Opinion: why we

need a centralized repository for isotopic data. *Proceedings of the National Academy of Sciences* **114**, 2997–3001. doi: 10.1073/pnas.1701742114

Pauls, S. U., Alp, M., Bálint, M., Bernabò, P., Čiampor, F., Čiamporová-Zaťovičová, Z., Finn, D. S., Kohout, J., Leese, F., Lencioni, V., Paz-Vinas, I., Monaghan, M. T. (2014). Integrating molecular tools into freshwater ecology: developments and opportunities. *Freshwater Biology* **59**, 1559–1576. doi.org/10.1111/fwb.12381

Peng, B., & Kimmal, M. (2005). simuPOP: a forward-time population genetics simulation environment. *Bioinformatics* **21**, 3686–3687.

Pine, W. E., Pollock, K. H., Hightower, J. E., Kwak, T. J., Rice, J. A. (2003) A review of tagging methods for estimating fish population size and components of mortality. *Fisheries* **28**, 10–3, doi: 10.1577/1548-8446(2003)28

Richter, K. K., Wilson, J., Jones, A. K. G., Buckley, M., van Doorn, N. & Collins, M. J. (2011). Fish 'n chips: ZooMS peptide mass fingerprinting in a 96 well plate format to identify fish bone fragments. *Journal of Archaeological Science* **38**, 1502–1510.

Ritchie, M. D., Holzinger, E. R., Li, R., Pendergrass, S. A. & Kim, D. (2015). Methods of integrating data to uncover genotype-phenotype interactions. *Nature Reviews Genetics* **16**, 85–97. doi:10.1038/nrg3868

Rodríguez-Ezpeleta, N., Bradbury, I. R., Mendibil, I., Álvarez, P., Cotano, U. & Irigoien, X. (2016). Population structure of Atlantic mackerel inferred from RAD-seq-derived SNP markers: effects of sequence clustering parameters and hierarchical SNP selection. *Molecular Ecology Resources* **16**, 991–1001. doi: 10.1111/1755-0998.12518

Rogers, A., Blanchard, J.L. & Mumby, P.J. (2014) Vulnerability of coral reef fisheries to a loss of structural complexity. *Current Biology* **24**, 1000–1005.

- Salipante, S. J., Kawashima, T., Rosenthal, C., Hoogestraat, D. R., Cummings, L. A., Sengupta, D. J., Harkins, T. T., Cookson, B. T. & Hoffman, N. G. (2014). Performance comparison of Illumina and ion torrent next-generation sequencing platforms for 16S rRNA-based bacterial community profiling. *Applied and Environmental Microbiology* **80**, 7583–7591. doi: 10.1128/AEM.02206-14
- Schrider, D. R. & Kern, A. D. (2014). Discovering functional DNA elements using population genomic information: A proof of concept using human mtDNA. *Genome Biology and Evolution* **6**, 1542–1548. doi.org/10.1093/gbe/evu116
- Serafy J. E., Shideler G. S., Araújo R. J., Nagelkerken I. (2015). Mangroves enhance reef fish abundance at the Caribbean regional scale. *PLoS ONE* **10**, e0142022. doi.org/10.1371/journal.pone.0142022
- Sevellec, M., Pavey, S. A., Boutin, S., Filteau, M., Derome, N. & Bernatchez, L. (2014). Microbiome investigation in the ecological speciation context of lake whitefish (*Coregonus clupeaformis*) using next-generation sequencing. *Journal of Evolutionary Biology* **27**, 1029–1046. doi: 10.1111/jeb.12374
- Silvertown, J. (2009). A new dawn for citizen science. *Trends in Ecology & Evolution*, **24**, 467–471. doi.org/10.1016/j.tree.2009.03.017
- Shin, Y.-J., Shannon, L.J., Bundy, A., Coll, M., Aydin, K., Bez, N., Blanchard, J.L., Borges, M.d.F., Diallo, I., Diaz, E., Heymans, J.J., Hill, L., Johannesen, E., Jouffre, D., Kifani, S., Labrosse, P., Link, J.S., Mackinson, S., Masski, H., Möllmann, C., Neira, S., Ojaveer, H., Abdallahi, K. O. M., Perry, I., Thiao, D., Yemane, D. & Cury, P. M. (2010). Using indicators for evaluating, comparing and communicating the ecological status of exploited marine ecosystems. 2. Setting the scene. *ICES Journal of Marine Science* **67**, 692–716.

968 Shogren, A. J., Tank, J. L., Andruszkiewicz, E., Olds, B., Mahon, A. R., Jerde, C. L. &  
 969 Bolster, D. (2017). Controls on eDNA movement in streams: transport, retention and  
 970 resuspension. *Scientific Reports* **7**, 5065. doi: 10.1038/s41598-017-05223-1  
 971 Sigsgaard, E. E., Nielsen, I. B., Bach, S. S., Lorenzen, E. D., Robinson, D. P., Knudsen, S.  
 972 W., Pedersen, M. W., Al Jaidah, M., Orlando, L., Willerslev, E. & Møller, P. R.  
 973 (2016). Population characteristics of a large whale shark aggregation inferred from  
 974 seawater environmental DNA. *Nature Ecology & Evolution* **1**,0004.  
 975 Smith, N. S. & Zeller, D. K. (2015) Unreported catch and tourist demand on local fisheries of  
 976 small island states: these case of The Bahamas, 1950-2010. *Fishery Bulletin* **114**,  
 977 117–131. doi: 10.7755FB.114.1.10  
 978 Song, W., Li, L., Huang, H., Jiang, K., Zhang, F., Chen, X., Zhao, M. & Ma, L. (2016). The  
 979 gut microbial community of antarctic fish detected by 16S rRNA gene sequence  
 980 analysis. *BioMedical Research International* Article ID 3241529. doi:  
 981 10.1155/2016/3241529  
 982 Sousa, L. L., Xavier, R., Costa, V., Humphries, N. E., Trueman, C., Rosa, R., Sims, D.W. &  
 983 Queiroz, N. (2016). DNA barcoding identifies a cosmopolitan diet in the ocean  
 984 sunfish. *Scientific Reports*, **6**, 1–9.  
 985 Stallings, C.D. (2009). Fishery-independent data reveal negative effect of human population  
 986 density on Caribbean predatory fish communities. *PLoS ONE* **4**, e5333.  
 987 doi.org/10.1371/journal.pone.0005333  
 988 Star, B., Nederbragt, A. J., Jentoft, S., Grimholt, U., Malmstrom, M., Greger, T. F., Rounge,  
 989 T. B., Paulsen, J., Solbakken, M. H., Sharma, A., Wetten, O. F., Lanzén, A., Winer,  
 990 R., Knight, J., Vogel, J.-H., Aken, B., Andersen, Ø., Lagesen, K., Tooming-  
 991 Klunderud, A., Edvardsen, R. B., Tina, K. G., Espelund, M., Nepal, C., Previti, C.,  
 992 Karlsen, B. O., Moum, T., Skage, M., Berg, P. R., Gjølén, T., Kuhl, H., Thorsen, J.,

993 Malde, K., Reinhardt, R., Du, L., Johansen, S. D., Searle, S. Lien, S., Nilsen, F.,  
 994 Jonassen, I., Omholt, S. W., Stenseth, N. C. & Jakobsen, K. S.(2011). The genome  
 995 sequence of Atlantic cod reveals a unique immune system. *Nature* **477**, 207–210.  
 996 Stark, A., Lin, M. F., Kheradpour, P., Pedersen, J. S., Parts, L., Carlson, J. W., Crosby, M.  
 997 A., Rasmussen, M. D., Roy, S., Deoras, A. N., Ruby, J. G., Brennecke, J., Harvard  
 998 FlyBase curators, Berkeley Drosophila Genome Project, Hodges, E., Hinrichs, A.  
 999 S., Caspi, A., Paten, B., Park, S.-W., Han, M. V., Maeder M. L., Polansky, B. J.,  
 1000 Robson, B. E., Aerts, S., van Helden, J., Hassan, B., Gilbert, D. G., Eastman, D.  
 1001 A., Rice, M., Weir, M., Hahn, M. W., Park, Y., Dewey, C. N., Pachter, L., Kent,  
 1002 W. J., Haussler, D., Lai, E. C., Bartel, D. P., Hannon, G. J., Kaufman, T. C., Eisen,  
 1003 M. B., Clark, A. G., Smith, D., Celniker, S. E, Gelbart, W. & Kellis, M. (2007).  
 1004 Discovery of functional elements in 12 Drosophila genomes using evolutionary  
 1005 signatures. *Nature* **450**, 219–232. doi.org/10.1038/nature06340  
 1006 Stuart-Smith, R. D., Bates, A. E., Lefcheck, J. S., Duffy, J. E., Baker, S. C., Thomson, R. J.,  
 1007 Stuart-Smith, J. F., Hill, N. A., Kininmonth, S. J., Airoidi, L., Becerro, M. A.,  
 1008 Campbell, S J., Dawson, T. P., Navarrete, S. A., Soler, G. A., Strain, E. M. A.,  
 1009 Willis, T. J. & Edgar, G. J. (2013) Integrating abundance and functional traits  
 1010 reveals new global hotspots of fish diversity. *Nature* **501**, 539–542.  
 1011 Tarnecki, A. M., Burgos, F. A., Ray, C. L. & Arias, C.R. (2017). Fish intestinal microbiome:  
 1012 diversity and symbiosis unravelled by metagenomics. *Journal of Applied*  
 1013 *Microbiology* **123**. doi: 10.1111/jam.13415  
 1014 Thomsen, P.F., Kielgast, J., Iversen, L. L., Møller, P. R., Rasmussen, M. & Willerslev, E.  
 1015 (2012). Detection of a diverse marine fish fauna using environmental DNA from  
 1016 seawater samples. *PLoS One* **7**, e41732.

1017 Thurstan, R. H., Brockington, S. & Roberts, C. M. (2010). The effects of 118 years of  
1018 industrial fishing on UK bottom trawl fisheries. *Nature Communications*, **1**, 15.

1019 Thurstan, R. H. & Roberts, C. M. (2010). Ecological meltdown in the Firth of Clyde,  
1020 Scotland: two centuries of change in a coastal marine ecosystem. *PloS ONE*, **5**,  
1021 e11767.

1022 Trueman, C. N., MacKenzie, K. M. & Palmer, M. R. (2012) Identifying migrations in marine  
1023 fishes through stable-isotope analysis. *Journal of Fish Biology* **81**, 826–847.

1024 Turner, C. R., Barnes, M. A., Xu, C. C., Jones, S. E., Jerde, C. L. & Lodge, D. M. (2014).  
1025 Particle size distribution and optimal capture of aqueous microbial eDNA. *Methods*  
1026 *in Ecology and Evolution* **5**, 676–684.

1027 Uchii, K., Doi, H. & Minamoto, T. (2016). A novel environmental DNA approach to quantify  
1028 the cryptic invasion of non- native genotypes. *Molecular Ecology Resources* **16**,  
1029 415–422.

1030 Ulibarri, R. M., Bonar, S. A., Rees, C., Amberg, J., Ladell, B. & Jackson, C. (2017).  
1031 comparing efficiency of American Fisheries Society standard snorkeling techniques  
1032 to environmental DNA sampling techniques. *North American Journal of Fisheries*  
1033 *Management* **37**, 644–651.

1034 Valentini, A., Taberlet, P., Miaud, C., Civade, R., Herder, J., Thomsen, P. F., Bellemain, E.,  
1035 Besnard, A., Coissac, E., Boyer, F. & Gaboriaud, C. (2016). Next- generation  
1036 monitoring of aquatic biodiversity using environmental DNA metabarcoding.  
1037 *Molecular Ecology* **25**, 929–942.

1038 Van Neer, W. & Ervynck, A. (2010). Defining “natural” fish communities for fishery  
1039 management purposes: biological, historical and archaeological approaches. In  
1040 *Ecologies and Economies in Medieval and Early Modern Europe* (Bruce, S. G., ed.),  
1041 pp. 188–218. Leiden: Brill.



1042 Vellend, M. (2003). Island biogeography of genes and species. *The American Naturalist*, **162**,  
1043 358–365.

1044 Vellend, M., Lajoie, G., Bourret, A., Múrria, C., Kembel, S. W. and Garant, D. (2014),  
1045 Drawing ecological inferences from coincident patterns of population and  
1046 community-level biodiversity. *Molecular Ecology*, **23**, 2890–2901.  
1047 doi: 10.1111/mec.12756

1048 Voegeli, F. A., Smale, M. J., Webber, D. M., Andrade, Y. & O'Dor, R. K. (2001). Ultrasonic  
1049 telemetry, tracking and automated monitoring technology for sharks. *Environmental*  
1050 *Biology of Fishes* **60**, 267–281.

1051 Weijerman, M., Fulton, E. A., Kaplan, I. C., Gorton, R., Leemans, R., Mooij, W. M. &  
1052 Brainard, R. E. (2015) An integrated coral reef ecosystem model to support resource  
1053 management under a changing climate. *PLoS ONE*, **10**, e0144165.

1054 Wheat, R. E., Wang, Y., Byrnes, J. E. & Ranganathan, J. (2013). Raising money for scientific  
1055 research through crowdfunding. *Trends in Ecology & Evolution* **28**, 71–2. doi:  
1056 10.1016/j.tree.2012.11.001

1057 Wheeler, A. & Jones, A. K. G. (1989). *Fishes. Cambridge Manuals in Archaeology*.  
1058 Cambridge: Cambridge University Press.

1059 Whoriskey, F. G. (2015). The ocean tracking network: a global partnership uses electronic  
1060 tagging technologies to track the movements of aquatic animals, answer science  
1061 questions, stimulate new technology development and assist with sustainable  
1062 development of the ocean. *Development and International Law* **47**, 221–232.

1063 Wilcox, T. M., McKelvey, K. S., Young, M. K., Lowe, W. H. & Schwartz, M. K. (2015).  
1064 Environmental DNA particle size distribution from Brook Trout (*Salvelinus*  
1065 *fontinalis*). *Conservation Genetics Resources* **7**, 639–641.

- Worm, B., Hilborn, R., Baum, J. K., Branch, T. A., Collie, J. S., Costello, C., Fogarty, M. J.,  
Fulton, E. A., Hutchings, J. A., Jennings, S., Jensen, O. P., Lotze, H. K., Mace, P.  
M., McClanahan, T. R., Minto, C., Palumbi, S. R., Parma, A. M., Ricard, D.,  
Rosenberg, A. A., Watson, R. & Zeller, D. (2009). Rebuilding global fisheries.  
*Science*, **325**, 578–585. doi.org/10.1126/science.1173146
- Xia, J. H., Lin, G., Fu, G. H., Wan, Z. Y., Lee, M., Wang, L., Liu, X. J. & Yue, G. H. (2014).  
The intestinal microbiome of fish under starvation. *BMC Genomics* **15**, 266. doi:  
10.1186/1471-2164-15-266
- Yamamoto, S., Minami, K., Fukaya, K., Takahashi, K., Sawada, H., Murakami, H., Tsuji, S.,  
Hashizume, H., Kubonaga, S., Horiuchi, T. & Hongo, M. (2016). Environmental  
DNA as a ‘snapshot’ of fish distribution: a case study of Japanese jack mackerel in  
Maizuru Bay, Sea of Japan. *PloS ONE* **11**, e0149786.
- Zeller, D., Harper, S., Zylich, K. and Pauly, D. (2015). Synthesis of underreported small-  
scale fisheries catch in Pacific island waters. *Coral Reefs*, **34**, 25–39.

### Electronic References

- Callou, C. (2009). Inventaires archéozoologiques et archéobotaniques de France (Inventaire  
National Du Patrimoine Naturel). *Muséum National d'Histoire Naturelle*. Available  
at [inpn.mnhn.fr/espece/jeudonnees/3471?lg=en](http://inpn.mnhn.fr/espece/jeudonnees/3471?lg=en)
- Davis, R., Baumgartner, M., Comeau, A., Cunningham, D., Davies, K., Furlong, A., Johnson,  
H., L'Orsa, S., Ross, T., Taggart, C. & Whoriskey, F. (2016). Tracking whales on the  
Scotian Shelf using passive acoustic monitoring on ocean gliders. *Oceans* 2016.  
Piscataway, NJ: Institute of Electrical and Electronics Engineers Available at  
[www.ieeeexplore.ieee.org/document/7761461/authors](http://www.ieeeexplore.ieee.org/document/7761461/authors)

1091 Froese, R. & Pauly, D. (Eds) (2017). FishBase. Available at [www.fishbase.org](http://www.fishbase.org).

1092 NASCO (2008) Salmon at Sea – SALSEA – An International Cooperative Research  
 1093 Programme on Salmon at Sea. *NASCO Report: SAL 04(5)*. Edinburgh: North  
 1094 Atlantic Salmon Conservation Organization.  
 1095 [www.nasco.int/sas/pdf/archive/other\\_reports/salsea\\_programme.pdf](http://www.nasco.int/sas/pdf/archive/other_reports/salsea_programme.pdf) (accessed 31st  
 1096 October 2017).

1097 OSPAR Convention. (2013). An assessment of the Ecological coherence of the OSPAR  
 1098 Network of Marine Protected Areas in 2012. London: OSPAR Commission.  
 1099 Available at  
 1100 [www.ospar.org/about/publications?q=An+assessment+of+the+Ecological+coherenc](http://www.ospar.org/about/publications?q=An+assessment+of+the+Ecological+coherenc)  
 1101 [e+of+the+OSPAR+Network+of+Marine+Protected+Areas+in+2012&a=&y=2013&](http://www.ospar.org/about/publications?q=An+assessment+of+the+Ecological+coherenc)  
 1102 [s=](http://www.ospar.org/about/publications?q=An+assessment+of+the+Ecological+coherenc)  
 1103  
 1104

TABLE 1. Summary of popular current and emerging methods used for fish surveys along with the associated advantages and limitations of each method.

Method	Advantages	Limitations	Personnel requirement	Health & safety consideration
Electric fishing	Can be used in flowing and still water, amongst macrophytes and obstructions; relatively unselective; can be used quantitatively	Inefficient in water > 1–1.5m depth or in wide reaches; limited by water and bed conductivity; can be harmful to sensitive fish species and life stages; invasive	Significant to high	High
Seine netting	Can be used quantitatively; efficiency well-understood; relatively unselective	Limited effectiveness in very deep or very shallow water; limited effectiveness where there are macrophytes, obstructions, or soft sediment; restricted to use in low velocity water bodies; invasive	High	Significant
Trawling	Large areas of deep water can be surveyed efficiently	Restricted to use in relatively open continuous stretches of water of > 2m depth; cannot be used where there are dense growths of macrophytes, very variable bed profiles or large debris; requires	High	High

		sizeable boats and launching facilities; invasive		
Gill netting	Can be used in a wide variety of environments amongst debris and macrophytes, in almost any depth	Invasive–destructive; limited ability to assess absolute fish abundance	Significant	Significant
Hydroacoustics	Huge expanses of water can be surveyed efficiently; non-invasive; quantitative abundance estimates possible	Limited effectiveness in turbulent environments; can only sample relatively open water so unsuitable to use for sampling in marginal habitats; lacks capacity to differentiate between species; cannot assess age, condition and health of fish	Significant	Significant
Fyke netting and trapping	Can be deployed in a variety of Environments; can be effective for some species difficult to sample by other methods	Very species and size-selective; limited ability to assess absolute fish abundance	Significant	Significant
Fry surveys: micromesh seine, handnet & traps	Focuses on margins of rivers and lakes, therefore less resource intensive, simple equipment only;	Only assesses juvenile populations; invasive as very young fish are unlikely to survive	Significant	Significant

	assesses a key life stage; relatively unselective	capture		
Fish counters & fixed traps: sometimes accompanied by cameras (video recorder)	Good for assessing highly mobile fish with relatively predictable migration patterns	Resource intensive, high capital costs, maintenance; quantitative assessment for migratory species only; often only operational under certain environmental conditions	High	Significant
Rod-and-line	Adaptable, can be deployed almost anywhere; amenable to volunteers and citizen-science participation	Very effort-dependent (quantity and quality); strongly influenced by conditions; very selective for species and size of fish; limited capability to assess absolute fish abundance; very noisy data	High	Low
Commercial catch monitoring	Enables large volumes of data collected over large spatial and temporal scales; relatively inexpensive as fish are being caught anyway	Can only happen where commercial fisheries exist; little control over changes in effort and methodology, which are driven by market forces; strongly influenced by conditions	Low	Low

Visual surveys: snorkelling, counting from the bank	Relatively non-invasive; enables observation of fish in their surroundings	Only applicable in high water clarity and over short ranges; mostly applicable to species with distinct individual home range, typically associated with physical habitat features	Moderate	Significant to high
Methods under development				
eDNA: single-target and meta barcoding	Very adaptable, deployable anywhere; non-invasive; non- selective; low field manpower requirement	Currently can only establish fish presence and abundance of species relative to each other, absolute abundance remains a challenge; cannot assess age, size, condition or health; uncertainty around the source of eDNA in lotic environments; high laboratory time requirement	Significant	Significant
DIDSON <sup>*</sup> –ARIS <sup>#</sup> high resolution sonar	Can be used in turbid water, amongst obstructions; can be used in a variety of depths and flows except very turbulent water; enables visualisation of target	Mobile deployment currently challenging; limited ability to assess whole water-body abundance; limited species identification capability; high	Significant	Significant

---

fish, species identification;	data-processing requirement;
quantitative estimates possible;	cannot assess age, condition
species (some) and size of fish can	and health of fish
be identified; observations of fish	
behaviour possible; non-invasive	

---

\*, Dual-frequency IDentification SONar; #, Adaptive Resolution Imaging Sonar.